

Successful Treatment of Pregnant and Postpartum Women With Severe COVID-19 Associated Acute Respiratory Distress Syndrome With Extracorporeal Membrane Oxygenation

JAIRO H. BARRANTES¹,* JAMEL ORTOLEVA,[†] ERIKA R. O'NEIL,[‡] ERIK E. SUAREZ,[§] SHARON BETH LARSON[¶], ANIKET S. RALI,^{*} CARA AGERSTRAND,^{||} LORENZO GRAZIOLI^{**}, SUBHASIS CHATTERJEE^{††} AND MARC ANDERS^{‡‡}

Abstract: There are limited data on the use of extracorporeal membrane oxygenation (ECMO) for pregnant and peripartum women with COVID-19 associated acute respiratory distress syndrome (ARDS). Pregnant women may exhibit more severe infections with COVID-19, requiring intensive care. We supported nine pregnant or peripartum women with COVID-19 ARDS with ECMO, all surviving and suffering no major complications from ECMO. Our case series demonstrates high-maternal survival rates with ECMO support in the management of COVID-19 associated severe ARDS, highlighting that these pregnant and postpartum patients should be supported with ECMO during this pandemic. *ASAIO Journal* 2021; 67;132–136

Key Words: ECMO, extracorporeal membrane oxygenation, COVID, COVID-19, ARDS, pregnancy, peripartum

The pandemic associated with coronavirus SARS-CoV-2 and related clinical disease, COVID-19, has affected over 38 million people worldwide with over one million deaths.¹ The hormonal, cardiovascular physiology, and immunomodulatory changes during pregnancy increase susceptibility to respiratory infections and may predispose to severe presentations of the disease.^{2,3} Additionally, reports of severe COVID-19 infections in pregnant and peripartum women and the fetal effects are emerging.^{2,4–7}

Extracorporeal membrane oxygenation (ECMO) is an invasive support strategy for cardiac, respiratory, or combined

cardiorespiratory failure when conventional treatment options have failed. ECMO has been successfully deployed for the management of critical illness in pregnant and postpartum patients, including during the previous pandemic.^{8–10} The use of ECMO for acute respiratory distress syndrome (ARDS) during the H1N1 pandemic saved many maternal and fetal lives, however, few studies report the use of pregnant and postpartum ECMO during this pandemic.^{4–8,11}

Against this background, we present an international case series of pregnant and peripartum patients managed with ECMO for COVID-19 induced ARDS, with maternal data and data on fetuses and neonates as relevant.

Methods

Pregnant and postpartum patients with a polymerase chain reaction confirmed for SARS-CoV-2 infection supported with ECMO were identified by the collaborating institutions from February until September 2020. Descriptive statistical methods included median (minimum-maximum range) and frequency, n (%). Demographics, maternal pre-ECMO, ECMO characteristics, and neonatal outcomes were described. Adverse events during the ECMO course were also identified. Individual IRB approval was obtained by the collaborating institutions.

Results

Our cohort includes nine patients with median age 30 years (range 22–43 years), five of whom were within 48 hours postpartum, two peripartum, and two pregnant at the time of ECMO initiation. All but two patients reported respiratory symptoms of COVID-19 during the third trimester. All patients had severe ARDS with a median PaO₂/FiO₂ (PF ratio) of 62 mm Hg (54–100 mmHg) managed with invasive mechanical ventilation, 5 (56%) with inhaled epoprostenol, and 6 (67%) with prone positioning. Specific COVID-19 therapies before ECMO included remdesivir 4 (44%), ribavirin/lopinavir 2 (22%), convalescent plasma therapy 4 (44%), hydroxychloroquine 4 (44%), azithromycin 6 (67%), anticytokine 2 (22%), and glucocorticoids 5 (56%). Table 1 summarizes maternal characteristics, including comorbidities and COVID-19-related symptoms and therapies.

The median RESP score was 4 [–1 to 6] before ECMO cannulation.¹² All patients received venovenous ECMO support, one concurrently with an intra-aortic balloon pump due to elevated left ventricular end-diastolic pressure and signs of biventricular dysfunction. ECMO cannulation strategies varied with three patients undergoing femoral-femoral, three with femoral-internal jugular configuration, and three with

From the *Division of Pulmonary, Critical Care and Sleep Medicine, Baylor College of Medicine, Houston, Texas; †Department of Anesthesiology and Perioperative Medicine, Tufts Medical Center, Boston, Massachusetts; ‡Department of Pediatrics (Critical Care), Texas Children's Hospital, Baylor College of Medicine, Houston, Texas; §Department of Cardiovascular Surgery, DeBakey Heart and Vascular Center, Houston Methodist, Houston, Texas; ¶Department of Surgery and the Division of Cardiothoracic Surgery, University of Iowa, Iowa City, Iowa; ||Division of Pulmonary, Allergy and Critical Care Medicine, Columbia University College of Physicians and Surgeons/NewYork-Presbyterian Hospital, New York, New York; **Department of Anesthesia and Intensive Care, ASST Papa Giovanni, Bergamo, Italy; and ††Department of Surgery, Baylor St Luke's Medical Center, Baylor College of Medicine, Houston, Texas.

Submitted for consideration October 2020; accepted for publication in revised form November 2020.

Disclosure: The authors have no funding and conflicts of interest to report.

Correspondence: Jairo H. Barrantes, Division of Pulmonary, Critical Care and Sleep Medicine, Baylor College of Medicine, Houston, TX. Email: jairo.barrantesperez@bcm.edu.

Copyright © ASAIO 2020

DOI: 10.1097/MAT.0000000000001357

Table 1. Maternal Characteristics

	patient 1	patient 2	patient 3	patient 4	patient 5	patient 6	patient 7	patient 8	patient 9
Center	A	A	B	C	D	E	F	B	C
Maternal factors									
Age (years)	28	30	22	27	34	30	30	36	43
Race	Hispanic	Hispanic	Caucasian	African-American	Declined	Hispanic	Indian	African-American	Caucasian
Weight (kg)	90.7	100.0	70.0	63.2	72.0	88.5	70.0	94.5	117.1
Comorbidities	Obesity	Obesity SLE	None	None	None	Obesity Asthma	None	Obesity Diverticulitis	Obesity CAD HTN
Pregnancy-related comorbidities	None	Hypothyroidism	Obesity	Tonic-clonic seizures	None	None	None	None	Placenta previa Succenturiate placental lobe
Maternal COVID-19									
Symptom onset (days)	7	6	8	4	4	7	#	17	6
COVID-19 diagnosis	PCR	PCR	PCR	PCR	PCR	PCR	PCR	PCR	PCR
CRP (mg/mL)	12.1	15.5	6.7	#	28.94	69.1	20.9	169.5	#
White-cell count ($\times 10^{-3}/\text{mm}^3$)	7.45	7.8	8.16	4.96	41.0	7.4	16.6	26.8	12.9
LDH (U/L)	221	486	1,100	444	664	296	459	834	#
Troponin (ng/mL)	<0.006	<0.006	6.5	<0.03	0.059	#	0.021	0.083	#
Ferritin (ng/mL)	77	429	72	77.7	588	72	#	#	#
D-Dimer (mcg/mL)	1.34	4.63	15.25	2.20	2.38	0.502	#	3.91	12.1
Fibrinogen (mg/dL)	586	537	447	540	765	#	596	1,266	640
Echo ejection fraction (EF)	60%	60%, RV enlarged	40-45%	Normal EF	Normal EF	Normal EF	Normal EF	Normal EF	Normal EF
Invasive ventilation	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
PEEP/plateau (cm H ₂ O)	12 / 30	14 / 30	14 / 31	16 / 29	10 / 28	19 / 28	15 / 34	16 / 30	16 / 30
PF ratio prior cannulation	56	55	69	66	100	66	60	54	62
Inhaled nitric oxide	No	No	No	No	No	No	No	No	No
Inhaled prostacyclin	Yes	Yes	Yes	No	No	Yes	No	Yes	No
Prone positioning	Yes	Yes	Yes	No	Yes	No	Yes	Yes	No
COVID-19 targeted therapy before ECMO	1. HCL 2. Azithromycin 3. Tocilizumab 4. Convalescent plasma 5. Remdesivir	1. HCL 2. Azithromycin 3. Convalescent plasma 4. Tocilizumab 5. Ribavirin	1. HCL 2. Azithromycin 3. Placebo arm – Remdesivir	Convalescent plasma	1. Azithromycin 2. Remdesivir	1. Azithromycin 2. Convalescent plasma 3. Remdesivir	1. HCL 2. Lopinavir	1. Remdesivir 2. Azithromycin 3. Dexamethasone	Dexamethasone
Systemic steroids	Yes	Yes	Yes	No	No	No	No	Yes	Yes
Gravida/Para	4/2	2/1	1/1	3/1	1/0	4/3	1/0	5/4	1/0
Gestational age	30+3	35+2	29	23	33	30	37+3	25+2	32+6
Delivery	CS	CS	CS	#	CS	CS	CS	CS	CS

#Data not available.

CAD, coronary artery disease; CS, cesarean section; HCL, hydroxychloroquine; HTN, arterial hypertension; PCR, Polymerase chain reaction; PEEP, positive end expiratory pressure; PF ratio, $\text{PaO}_2/\text{FiO}_2$; RV, right ventricle; SLE, systemic lupus erythematosus.

single-site dual lumen right internal jugular cannulation. Anticoagulation was initiated with unfractionated heparin bolus in 8 (89%) patients (with one not receiving a heparin bolus due to bleeding concerns), with six patients maintained on unfractionated heparin infusion, but three patients changed to bivalirudin targeting institutional thresholds for anti-Xa, partial thromboplastin time, or thromboelastography. Two patients developed thrombotic complications of the membrane oxygenator during ECMO support, managed with oxygenator exchange. One patient developed minor vaginal bleeding during ECMO, and one patient underwent surgical exploration after ECMO decannulation for surgical site bleeding. The median ECMO duration for the first ECMO run was 10 days (range 6–57 days). Our most obese patient (patient 9C) was supported with ECMO flows up to 6.5 L/min without complications and minimal recirculation. Targeted COVID-19 treatment during ECMO included antiviral therapy (remdesivir, ribavirin), and immunomodulatory therapy (convalescent plasma therapy, tocilizumab, Cytosorb). All patients were successfully decannulated from ECMO; however, one patient (9C) required a second 10 day VV-ECMO run (femoral-femoral) postpartum due to worsening respiratory failure. All patients were discharged from the hospital alive, except for two patients still admitted. Further details of the ECMO support are outlined in Table 2.

The two patients with ECMO initiation during pregnancy were electively cannulated in the operating room. While supine positioned, cannulation was directed by fluoroscopy and transesophageal echocardiography. Regular follow-up chest x-rays confirmed cannula positioning. Deliberately both patients were cannulated via single-site dual lumen right internal jugular cannula due to potential concern for compression of the uterus and the intraabdominal and pelvic vessels, potentially limiting uterine blood supply. No repositioning of the cannula was required during each of the ECMO runs. Both pregnant patients were able to perform physical therapy and ambulated. Both fetuses were monitored via cardiotocography and ultrasound during ECMO support.

All newborns were delivered via cesarean section and all but one were premature, less than 37 weeks gestation. Evaluation of the two fetuses for the two patients supported with ECMO during pregnancy demonstrated normal fetal growth and no overt fetal compromise. One patient (4C) gave birth 106 days post ECMO cannulation at term gestation, with good APGARs (7/9), the other patient (9C) delivered after completion of the first ECMO run. None of the followed-up newborns had laboratory or clinical evidence suggestive of COVID-19 (Table 3). Maternal breastmilk was not used for any of the newborns except in the one patient, who gave birth 106 days after ECMO decannulation. One newborn died before hospital discharge.

Discussion

This case series provides insights for pregnant patients supported with ECMO for COVID-19 ARDS. All patients were successfully weaned off ECMO and all except one neonate survived despite maternal distress and preterm delivery. ARDS management, COVID-19 related therapies, as well as ECMO support, did not result in an immediate negative impact on the mothers nor infants. Finally, none of the infants contracted

COVID-19 but the majority were premature and required intensive care.

The use of ECMO for pregnant women during the novel coronavirus pandemic can be challenging given the limited data.^{4–7} In general, however, pregnant women are younger, healthier, and have close medical follow-up when compared with the general population. This likely contributes to the high survival with the use of ECMO for other indications in peripartum women.^{8–10} ECMO for COVID-19 ARDS in peripartum patients is described in two case reports.^{5,6} Additionally, two European studies describe 1% of their peripartum patients with COVID-19 requiring ECMO support, however, no details on the patients, ECMO run, nor complications are given.^{4,7} Recently, Barbaro et al reported on the successful use of ECMO in COVID-19 in an international cohort study, where 22 patients out of 1,035 were pregnant.¹¹ Our case series highlights high survival with the support of ECMO for ARDS related to COVID-19 and provides details that may be useful to physicians caring for severely ill pregnant women.

Before the initiation of ECMO, ARDS management for pregnant women followed the clinical practice guidelines for mechanical ventilation in adult patients with ARDS.¹³ All patients were treated with high PEEP and low driving pressures, neuromuscular blockade, and 6/9 (67%) with prone positioning.¹⁴ Specific COVID-19 treatment options varied and included systemic glucocorticoids, antiviral, and other immunomodulatory therapeutics. As in the previous pandemic, once conventional intensive therapies were exhausted, ECMO support for ARDS was initiated for these women.⁸

Similar to the general population affected by COVID-19, pregnant women with comorbidities such as obesity, diabetes, and hypertension are at higher risk of serious infection with COVID-19, with 10% needing critical care.^{4,7} Comparable to the patient Fiore et al described the majority of our patients were obese.⁵ The overall most common complication on ECMO for peripartum patients remains bleeding.⁹ Two of our patients experienced minor bleeding but no major complications during ECMO, alike to the two cases reported.^{5,6} The most common complication reported in ECMO for COVID-19 ARDS was circuit change (15%), which mirrors our complication rate for circuit/oxygenator clotting (22%).¹¹ Circuit thrombosis may be more common as a result of the combination of pathophysiological alterations of hemostasis during pregnancy, and the anticoagulation protocols, which were the standard practice of each institution and not modified for pregnancy nor COVID-19.

Fortunately, vertical transmission of COVID-19 from mother to fetus is rarely reported.¹⁵ None of our infants contracted COVID-19. The majority of our infants were premature, similar to other reports.^{6,15} Not previously reported, the majority of these infants required admission to intensive care, 71% required mechanical ventilation, and one infant is still admitted, and receiving noninvasive positive pressure ventilation. Despite a perilous gestation, the majority of our infants survived.

We recognize that our data and interpretation are limited by the small sample size. However, during this unprecedented pandemic as management strategies and therapies are continually evolving, we felt it important to share the data to help physicians at the bedside.

Table 2. ECMO Characteristics

Patient/Center	1/A	2/A	3/B	4/C	5/D	6/E	7/F	8/B	9/C
Canulation timing	Postpartum	Postpartum	Postpartum	Pregnancy	Postpartum	At delivery	Postpartum	At delivery	Pregnancy
Canulation type	VV	VV	VV	VV	VV	VV	VV	VV	VV
PF ratio prior cannulation	56	55	+IABP (x5 days) 69	66	100	66	60	54	62
RESP score ⁵	5	3	4	0	6	5	4	4	-1
Canulation location	Fem (25F)→ Fem (21F)	Fem (19F)→ Fem (21F)	Fem (25F)→ Fem (21F)	RIJ-Dual Lumen (31F)	Fem (23F)→ RIJ (20F)	Fem (25F)→ RIJ (19F)	RIJ - Dual Lumen (31F)	Fem (29F)→ RIJ (24F)	RIJ - Dual Lumen (31F)
Anticoagulation on ECMO	Heparin, changed to Bivalirudin due circuit clotting with therapeutic heparin	Heparin, changed to Bivalirudin (suspicion of HIT)	Heparin	Heparin	Heparin	Heparin	Heparin	Heparin, changed to Bivalirudin (suspicion of HIT)	Heparin
Anticoagulation parameters	PTT (60–90s)	PTT (60–90s)	PTT (60–80s)/ Anti-Xa (0.25–0.3)/TEG (R) 2–3 baseline	PTT (60–90s)/ TEG (R) 2–3 baseline	PTT (60–80s)	Anti-Xa (0.2–0.4)	PTT/TEG (various aims)	PTT (60–80s)/ Anti-Xa (0.25–0.3)/TEG (R) 2–3 baseline	PTT (60–90s)/ TEG (R) 2–3 baseline
Complications on ECMO	1. Circuit clotted on Heparin 2. Transient AKI 3. Oxygenator changed for fibrin deposition	1. Inotropes used 2. Right sciatic nerve high-grade stretch injury with foot drop	1. Inotropes used 2. SIADH 3. CINMP	1. Vasodilators required 2. Hyperbilirubinemia	Tension pneumothorax before and on ECMO	1. Surgical site bleeding 2. Bacterial pneumonia 3. Oxygenator clotted	1. Cytosorb 2. Systemic steroids	1. Inotropes used 2. Vaginal bleeding 3. Infection	None
ECMO duration (days)	9	11	7	9	10	6	13	57	14
Weaned off ECMO	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
COVID-19 targeted therapy on ECMO	1. Remdesivir (started pre-ECMO) 2. Systemic steroids	Ribavirin	1. Placebo arm— Remdesivir after study unblinded started after decannulation	1. Remdesivir—continued from before ECMO 2. Convalescent plasma RCT	1. Remdesivir continued from before ECMO 2. Convalescent plasma	1. Remdesivir continued from before ECMO 2. Convalescent plasma	1. Cytosorb 2. Systemic steroids	1. Systemic steroids 2. Convalescent plasma	Systemic steroids
Post ECMO complications	Bilateral lower extremities DVT (former ECMO site)	Vaginal bleeding	None	None	None	1. Left common femoral vein DVT (former ECMO site) 2. Surgical site exploration for bleeding	GBS	None	Recannulation 15 days after decannulation, 2nd ECMO run 10 days in duration, vented via trach
Maternal survival to hospital discharge	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Pending	Pending

#Data not available.

AKI, acute kidney injury; CIPNM, critical illness polynuropathy and myopathy; Fem, femoral vein; GBS, Guillain-Barré syndrome; HIT, heparin-induced thrombocytopenia; RIJ, right internal jugular vein; SIADH, syndrome of inappropriate antidiuretic hormone.

Table 3. Newborn/Fetal Characteristics at Time of ECMO

Patient/Center	1/A	2/A	3/B	4/C	5/D	6/E	7/F	8/B	9/C
Fetal monitoring during ECMO (CTG/US/other/none)	Postpartum ECMO	Postpartum ECMO	Postpartum ECMO	CTG and US	Postpartum ECMO	Postpartum ECMO	Postpartum ECMO	Postpartum ECMO	CTG and US
APGAR (1/5 minutes)	7/8	8/9	1/3	*	*	2/2	6/7	1/4	1/3
Gestational age	30 + 3	35 + 2	29	23	33	30	37 + 3	25 + 2	32 + 6
Admission to NICU	Yes	*	Yes	*	*	Yes	Yes	Yes	Yes
COVID status	Negative	Negative	Negative	*	*	Negative	Negative	Negative	Negative
Congenital abnormalities	None	None	None	*	*	None	None	None	PFO
Mechanical ventilation	Yes	No	Yes	*	*	Yes	No	Yes	Yes
Vasoactives	No	No	Yes	*	*	No	No	Yes	No
Newborn survival to hospital discharge	Yes	Yes	No	Yes†	*	Yes	Yes	Pending	Yes

*Data not available (local IRB restricted use).

†Delivered 106 days post ECMO decannulation.

CTG, cardiotokogram; PFO, persistent foramen ovale; US, ultrasound.

Conclusion

Our case series demonstrates excellent maternal and neonatal survival rates and supports the successful use of respiratory ECMO in the management of COVID-19 associated severe ARDS in pregnant and postpartum patients at high volume ECMO centers. As COVID-19 continues to impact thousands of patients worldwide daily, and despite limited data and resources, pregnancy should not be considered a contraindication for ECMO support for COVID-19 ARDS.

References

- World Health Organization (WHO): *Coronavirus disease (COVID-19) dashboard*. 2020. Available at <https://covid19.who.int/>. Accessed October 11, 2020.
- Chen YH, Keller J, Wang IT, Lin CC, Heng-Ching L: Pneumonia and pregnancy outcomes: A nationwide population-based study. *Am J Obstet Gynecol* 207: e1–e7, 2012.
- Center for Disease Control and Prevention (CDC): *Data on COVID-19 during pregnancy*. 2020. Available at <https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/special-populations/pregnancy-data-on-covid-19.html>. Accessed October 11, 2020.
- Kayem G, Lecarpentier E, Deruelle P, et al: A snapshot of the Covid-19 pandemic among pregnant women in France. *J Gynecol Obstet Hum Reprod* 49: 101826, 2020.
- Fiore A, Piscitelli M, Adodo DK, et al: Successful use of extracorporeal membrane oxygenation postpartum as rescue therapy in a woman with COVID-19. *J Cardiothorac Vasc Anesth* 1–4, 2020.
- Hou L, Li M, Guo K, et al: First successful treatment of a COVID-19 pregnant woman with severe ARDS by combining early mechanical ventilation and ECMO. *Heart Lung* 50: 33–36, 2020.
- Knight M, Bunch K, Vousden N, et al: UK Obstetric Surveillance System SARS-CoV-2 Infection in Pregnancy Collaborative Group: Characteristics and outcomes of pregnant women admitted to hospital with confirmed SARS-CoV-2 infection in UK: National population based cohort study. *BMJ* 369: m2107, 2020.
- Nair P, Davies AR, Beca J, et al: Extracorporeal membrane oxygenation for severe ARDS in pregnant and postpartum women during the 2009 H1N1 pandemic. *Intensive Care Med* 37: 648–654, 2011.
- Naoum EE, Chalupka A, Haft J, et al: Extracorporeal life support in pregnancy: A systematic review. *J Am Heart Assoc* 9: e016072, 2020.
- Ramanathan K, Tan CS, Rycus P, et al: Extracorporeal membrane oxygenation in pregnancy: An analysis of the extracorporeal life support organization registry. *Crit Care Med* 48: 696–703, 2020.
- Barbaro RP, MacLaren G, Boonstra PS, et al: Extracorporeal Life Support Organization: Extracorporeal membrane oxygenation support in COVID-19: An international cohort study of the extracorporeal life support organization registry. *Lancet* 396: 1071–1078, 2020.
- Schmidt M, Bailey M, Sheldrake J, et al: Predicting survival after extracorporeal membrane oxygenation for severe acute respiratory failure. The Respiratory Extracorporeal Membrane Oxygenation Survival Prediction (RESP) score. *Am J Respir Crit Care Med* 189: 1374–1382, 2014.
- Fan E, Beitler JR, Brochard L, et al: COVID-19-associated acute respiratory distress syndrome: Is a different approach to management warranted? *Lancet Respir Med* 8: 816–821, 2020.
- Bikash RR, Trikha A: Prone position ventilation in pregnancy: Concerns and evidence. *J Obst Anaesth Crit Care* 8: 7–9, 2018.
- Dumitriu D, Emeruwa UN, Hanft E, et al: Outcomes of neonates born to mothers with severe acute respiratory syndrome coronavirus 2 infection at a large medical center in New York City [published online ahead of print October 12, 2020]. *JAMA Pediatr*: e204298, 2020. doi:10.1001/jamapediatrics.2020.4298.