

Original article

Title: Clinical characteristics, management, and short term outcome of neonates born to mothers with COVID-19 in a tertiary care hospital in India

Sushma Malik, MD¹*, Suchitra Surve, MD², Poonam Wade, MD¹, Santosh Kondekar, MD¹, Vishal Sawant, MD¹, Murtuja Shaikh, MD¹, Periyasamy Kuppusamy, PhD², Ramesh Bharmal, MD³, Smita D Mahale, PhD², Deepak N Modi, PhD², Rahul K Gajbhiye, MBBS, PhD²*

¹Department of Paediatrics & Neonatology, Topiwala National Medical College & BYL Nair Charitable Hospital, Mumbai, India, 400 008.

²ICMR-National Institute for Research in Reproductive Health, Mumbai, India, 400 012.

³Dean, Topiwala National Medical College & BYL Nair Charitable Hospital, Mumbai, India, 400 008.

***Corresponding author**

Dr Sushma Malik, MD
Professor and Head, Department of Paediatrics & Neonatology
Topiwala National Medical College &
BYL Nair Charitable Hospital, Mumbai, India, 400 008.

Email : sushmamalik@gmail.com

Telephone + 919819065322

Or

Dr Rahul K Gajbhiye, MBBS, PhD
Scientist D & DBT Wellcome India Alliance

Clinical & Public Health Intermediate Fellow,
ICMR-National Institute for Research in Reproductive Health,
Parel, Mumbai, 400012, INDIA
Telephone: +91 22 24192036
Email: gajbhiyer@nirrh.res.in

Word count : Abstract : 213, Manuscript : 2946

Lay summary: The current pandemic of COVID-19 has affected all the countries globally. However, the adverse impact of the pandemic is more seen in the low-income and middle-income countries (LMICs). Although there is evidence on the adverse impact of the SARS-CoV-2 on the health of mothers and neonates, the evidence is mainly from high-income countries. For reducing the mortality and morbidity due to COVID-19 in LMICs, there is a need to generate evidence from the LMICs. The present study is a part of the National Registry of pregnant women with COVID-19 in India (PregCovid registry). Our study demonstrates a higher risk of adverse outcomes such as neonatal sepsis and death in the SARS-CoV-2 infected as compared to the non-infected neonates. The study also showed the risk of SARS-CoV-2 infection in 6.3 % of neonates born to mothers with COVID-19.

Introduction

The emergence of the outbreak of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in December 2019 has caused a global pandemic resulting in potentially lethal coronavirus disease 2019 (COVID-19). More than 167 million COVID-19 cases and more than 3.4 million deaths have been reported due to COVID-19 as of 27th May 2021[1]. Though it affects all age groups, there is growing evidence of its impact on pregnant women and their newborns [2]. India continues to be amongst the top countries having increasing numbers of COVID-19 cases affecting nearly 27 million people with more than 0.3 million deaths as of 27th May 2021 [3]. India accounts for nearly one-fifth of the world's annual childbirths thereby suggesting possibly large numbers of neonates affected with SARS-CoV-2 infection.

Current evidence indicates no major adverse neonatal outcomes of that maternal COVID-19 and a low risk of vertical transmission [4-6]. However, most of the data is accumulated from high-income countries [6-9]. There is very little information about clinical presentations and management of neonates born to COVID-19 mothers in low-income and middle-income countries (LMICs). There is a need to document the same in different populations.

The possibility of the maternal-fetal transmission of the SARS-CoV-2 is high due to various reasons such as the expression of ACE-2 receptor in the placenta [10]. However, the lower intracellular response induced by ACE-2 in the alveolar epithelial cells and the presence of alpha and gamma chains in fetal hemoglobin may exhibit a protective response in newborns [11]. Despite this hypothesis, severe course, and adverse outcomes are reported in neonates [12] and the occurrence of vertical transmission is demonstrated in some of the studies [13-15]. However, early neonatal infection due to horizontal transmission and/or environmental contamination cannot be excluded in some of the studies. There is evidence of SARS-CoV-2 detection in the

immediate skin-to-skin contact in vigorous babies was encouraged. Mothers performed hand hygiene and wore a mask. Those babies who required resuscitation were immediately shifted under the pre-warmed radiant heat warmer in the designated newborn resuscitation room and were appropriately managed as per the standard NRP 2015 resuscitation guidelines with a minimum number of personnel attending (one doctor for low-risk cases and two for complicated cases). COVID-19 status did not alter the indications for intubation.

New-born testing strategies

The nasopharyngeal swab was sent for all babies for a real-time reverse-transcription–polymerase chain reaction (RT-PCR) assay for SARS-CoV-2. From 14th April-May 2020 the first swab was sent after 24 hours of birth and the second swab after 72 hours of birth as per guidelines [21]. With the change of national policy, from June 2020 - till 31st July 2020 only the first swab was sent at birth / within 12 hours of life. If the first swab was negative and the baby was asymptomatic no further testing was done. In cases where the first swab was positive, a repeat swab was collected 24-48 hours later and tested.

Place of care, feeding, and management policies

All stable newborns were kept in the postnatal ward with mothers irrespective of their COVID-19 status (Zero separation policy) [22,23]. All neonates born to COVID-19 positive mothers who were roomed-in with their mothers in post-natal neonates were initiated into early and continued exclusively breastfed as far as possible. Occasionally, in a few babies, pasteurized donor human milk from Human Milk Bank or acceptable animal milk from pasteurized dairy milk with standardized fat composition was given. All the newborns were managed as per guidelines by expert group consensus. The antibiotics were given only if indicated in clinically suspected sepsis or proven sepsis [22]. The antibiotics given were injectable cefotaxime and amikacin

SARS-CoV-2 testing in neonates

Thirty-three of the 523 neonates (6.3%) were positive for SARS-CoV-2 by RT-PCR while the others were negative during the stay in the hospital. We compared the presentation and the outcomes of these 33 SARS-CoV-2 positive and 490 SARS-CoV-2 negative neonates born to mothers with COVID-19. The details of the 33 neonates are shown in Table 1. Amongst 33 COVID-19 positive neonates, 29 newborns were positive in their first swab itself collected within 12-24h of birth. There were four neonates whose initial RT-PCR test was negative; however, their repeat test at 72 hours of life was positive. In most neonates, the repeat RT-PCR test done after 48-72h of the first test was negative except for four neonates where the repeat test was also positive. Six of the 33 SARS-CoV-2 positive neonates were symptomatic (24%) while the others were asymptomatic. The profile and course of newborns of the mothers with COVID-19 are presented in Figure 1.

Presentations and outcomes in SARS-CoV-2 positive and negative neonates

Out of 524 neonates, 53.6% (n=281) were males and 46.4% (n=243) were females (Table 2). The majority (95.2%, 499 out of 524) were born at full-term gestation and only 4.8% (25/524) were born as preterm. Amongst them, 88.5% (n=464) newborns were born appropriate for their gestation and 11.3% were born as small for their gestational age, one newborn was large for the gestational age. Nearly 76% new-borns had birth weight >2.5kg, 22.5% had 1.5 to 2.5kg and 1.1% had less than 1.5kg. Most of the babies (95.4%) cried immediately after birth. No significant differences were observed between maturity at gestation, birth weight, and sex of SARS-CoV-2 infected and non-infected babies.

Amongst 523 newborns, 19.9% (n=104) newborns develop complications such as fetal distress, hypoglycemia, hypocalcemia, meconium aspiration, sepsis, etc. during the neonatal period; 133

Page 9 of 31

Manuscripts submitted to Journal of Tropical Pediatrics

9

(26%) required NICU admission (Figure 2). Of these majority were for reasons like hyperbilirubinemia, prematurity and low birth weight, respiratory distress, and other congenital anomalies like cleft lip & cleft palate, hydronephrosis, occipital encephalocele, and congenital diaphragmatic hernia. Nearly 8% of newborns were admitted for social reasons (baby admitted for care because of mother’s illness). Out of 523 neonates, 97.1% (n=509) were either discharged or in a stable condition at the time of data extraction.

A comparison of the clinical presentations and complications is shown in Table 3. Sepsis was present in a greater proportion among SARS-COV-2 infected neonates as compared to uninfected neonates (12.1% vs 3.3%). The risk of sepsis was 4.09 folds (95% CI,1.28-13.00) higher in the neonates with SARS-CoV-2 infected group as compared to the non-infected group (p=0.031). Of the 33 SARS-CoV-2 infected newborns 3% had hypocalcemia, 6.1% had hypoglycemia and 15.2% had hyperbilirubinemia. The risk of hypocalcemia was 7.6 (95% CI, 0.67-86.34) folds and hypoglycemia was 4.4 (95% CI, 0.89-22.33) folds higher amongst SARS-CoV-2 infected neonates as compared to uninfected neonates. Similarly, breathing difficulties (respiratory distress) were also reported significantly higher among SARS-CoV-2 infected neonates (12.1%) than non-infected (3.9%) (p=0.05).

The risk of developing fever in SARS-CoV-2 infected neonates was higher (OR 5.01, 95% CI, 0.51-50.16) as compared to non-infected newborns, although this increase was not significant statistically (Table 4). Poor feeding was significantly more common among SARS-CoV-2 infected neonates (12.1%) as compared to SARS-CoV-2 non-infected (2.7%) neonates (p=0.017). The risk of having poor feeding was 5.06 (1.55-16.49) times and respiratory distress was 3.42 folds (95% CI, 1.09-10.71) higher in SARS-CoV-2 infected neonates as compared to

http://mc.manuscriptcentral.com/tropej

Downloaded from https://academic.oup.com/tropej/advance-article/doi/10.1093/tropej/fmab054/6296618 by guest on 12 June 2021

their uninfected counterparts. Amongst the cohort of 33 infected neonates, one newborn had a patchy shadow suggestive of pneumonia.

The comparison of management strategies used for infected and non-infected neonates of mothers with COVID-19 is presented in Table 5. Nine out of 33 SARS-CoV-2 infected neonates required intensive care for management, while 123/490 non-infected neonates required NICU admission (OR 1.1, 95 % CI, 0.50-2.45). The need for bag and mask, CPAP, and the ventilator was, 1.91, 6.00, and 3.67 times higher in SARS-CoV-2 infected neonates than non-infected, however, the use of CPAP was only statistically significant.

As shown, in Table 6, out of nine neonates who were SARS-CoV-2 infected required NICU admission, two neonates required intensive care because of birth asphyxia and diabetes in the mother whereas three neonates had hyperbilirubinemia and required phototherapy. The remaining four neonates had clinically suspected sepsis and amongst them, two neonates were preterm and one was small for gestational age. Hemoglobin was decreased in case 4 and case 7. The leucocyte or white blood counts were decreased in case no 8. Low platelet count was seen in cases 4, 7, and 8. None of the neonates had grossly deranged liver function tests or renal function tests. However, C reactive protein (CRP) was elevated in cases 4, 6, 7, and 8. Other markers of inflammation like Interleukin 6 (IL-6) were elevated in case 6 and case 8. Blood culture was normal in cases 2, 4, and 8 whereas it grew *Klebsiella Pneumonia* in cases 6 and case 7. Thus, an increased risk of sepsis was observed in neonates with SARS-CoV-2 infection.

There was a total of 13 neonatal deaths, of which three deaths occurred in SARS-CoV-2 infected neonates (9%) who required intensive care while 10 (3%) were reported in the SAR-CoV-2 negative group. The risk of neonatal death was higher in SARS-CoV-2 infected newborns [(OR (4.8, 95% CI1.25-18.36).

Discussion

We analyzed the clinical characteristics of neonates born to COVID-19 mothers and found that the rate of SARS-CoV-2 infection in neonates was 6.3%. Compared to their uninfected counterparts, these neonates have a higher chance of developing clinical complications and required intense medical management. There is also a high risk of neonatal mortality amongst the SARS-CoV-2 infected neonates. The results of the study showed no significant difference between the proportion of males and females as predisposing factors to COVID-19. This observation was similar to the study reported by Dong et al showing that sex did not affect the incidence or severity of COVID-19 [25]. In our cohort, nearly 4.8% of the neonates were born as preterm which is lesser than that reported previously which has suggested high rates of preterm birth and stillbirths in SARS-CoV-2 infected women [5,26,27]. Factors like the appropriate implementation of protocols by the hospital for management of COVID-19 mothers may cause reduced rates of preterm births, improved air quality, home care, and reduced stress due to lockdown may also contribute to such reduced preterm birth rates. However, increased rates of home deliveries due to reduced access to the healthcare system could lead to under-reporting. Nevertheless, most neonates born to COVID-19 mothers were generally healthy and did not require special medical attention. Most newborns were in stable condition without any adverse outcome reiterating favorable course reported in newborns with lower fatality rates [28,29].

Infection of the newborn with SARS-CoV-2 is a controversial topic. Large systematic reviews have reported the incidence of SARS-CoV-2 infection in nearly 3-7% of neonates born to COVID-19 mothers [2,26,31]. Corroborating this data, herein we observed that nearly 6% of neonates were positive for SARS-CoV-2. Since we did not analyze the amniotic fluid or placenta, it is hard to comment if this virus was vertically transmitted or acquired postnatally. As

per the current case definition of mother-to-child transmission, a neonate positive for SARS-CoV-2 at birth but not after 24 or 48h is a probable case of intrapartum transmission [32]. In our study as most of the neonates that were positive for SARS-CoV-2 at birth, the repeat swab collected at 48h was negative for the virus. Thus in most cases, the virus seems to be acquired intrapartum in our cohort.

In a systematic review of 105 neonates which reported 8.8% of positivity [28], the rate of premature was lower amongst the infected neonates. However, in our study, although the numbers of premature neonates were higher in the infected group as compared to uninfected controls, the difference was not statistically significant. Nevertheless, our analysis revealed that the SARS-CoV-2 infected neonates are at increased risk of neonatal complications as compared to their uninfected counterparts despite no difference in their maturity status or birthweight. This observation is intriguing and warrants investigation as the current notion believes that SARS-CoV-2 infection is rare and not deleterious to the newborn. Our study highlights that newborns with SARS-CoV-2 infection are more susceptible to the adverse clinical course as respiratory difficulties and poor feeding were more common among SARS-CoV-2 infected babies; mechanical ventilation and CPAP were more commonly required in SARS-CoV-2 infected babies. Whether respiratory distress and poor feeding can be predictors of severity of the outcome in infected newborns needs further investigation. It is important to note that all the nine SARS-CoV-2 infected neonates who required NICU admission had underlying medical conditions warranting NICU stay. Characteristically, low platelet count decreased white blood cells and elevated IL-6 were seen in neonates with clinical deterioration along with positive septic screen in terms of elevated CRP and blood culture in a couple of cases. This suggests that baseline investigations like complete blood count and CRP are essential and altered biochemical

parameters especially platelet count, CRP, IL6, and LDH are characteristically altered in SARS-CoV-2 infected neonates with severe adverse outcome including death similar to findings reported by McLaren et al [30].

An intriguing observation we made was a high risk of death amongst SARS-CoV-2 infected newborns as compared to their non-infected counterparts. These neonates were not just positive for SARS-CoV-2 but also developed sepsis. Previous studies have not reported deaths amongst the SARS-CoV-2 infected neonates [7,33], however, the sample size is small in most studies, and in many instances, it is unknown if the mother was also infected with COVID-19 [34]. Furthermore, most studies reported neonates in a wide age range, while in our study we analyzed the death rates within the early neonatal period. As maternal COVID-19 would have some pathophysiologic effects including that in the placenta [13,14], this may have affected the newborn making them more susceptible to adverse outcomes. We believe that maternal COVID-19 would compound the impact of SARS-CoV-2 on newborns and there is a need to address this angle. It will be of clinical relevance to follow up with the non-infected neonates and investigate their susceptibility to COVID-19 and other infections.

To summarize, this is probably the first large study describing the clinical profile of neonates born to SARS-CoV-2 infected mothers and its significance with associated morbidities, especially in the COVID-19 newborns. The strength of the study lies in the fact that this is a first detailed study of a large cohort of newborns of COVID-19 mothers investigated uniformly. The limitations include single-center data and lack of long-term follow-up of the neonates. In absence of the data from uninfected mothers, we are unable to compare the outcomes and comment if maternal COVID-19 is a major risk factor for neonatal health. Also in our cohort, most mothers were infected with SARS-CoV-2 near term and hence the impact of early trimester infections on

newborn outcomes can not be judged. Nevertheless, our study highlights the need for further research exploring the compounding health risk of a newborn in presence of maternal COVID-19.

Acknowledgments

The authors acknowledge the network of the National Registry of Pregnant women with COVID-19 in India (PregCovid Registry, CTRI/2020/05/025423). All faculties, resident doctors, and nurses in the department of Paediatrics and Obstetrics at TNMC & BYL Nair Hospital, Mumbai are sincerely acknowledged. The study is supported by an intramural grant of ICMR-NIRRH (ICMR-NIRRH/RA/1036/02-2021). RG is an awardee of the DBT Wellcome India alliance clinical and public health intermediate fellowship (Grant no. IA/CPHI/18/1/503933).

Funding: No funding.

References

1. WHO Coronavirus Disease (COVID-19) Dashboard n.d. <https://covid19.who.int> (27 May 2021, date last accessed).
2. Gajbhiye R, Modi D, Mahale S. Pregnancy outcomes, Newborn complications and Maternal-Fetal Transmission of SARS-CoV-2 in women with COVID-19: A systematic review of 441 cases. *MedRxiv* 2020:2020.04.11.20062356.
3. MoHFW | Home n.d. <https://www.mohfw.gov.in/> 27 May 2021, date last accessed).
4. Adhikari EH, Moreno W, Zofkie AC, et al. Pregnancy Outcomes Among Women With and Without Severe Acute Respiratory Syndrome Coronavirus 2 Infection. *JAMA Netw Open* 2020;3.
5. Oncel MY, Akın IM, Kanburoglu MK, et al. A multicenter study on epidemiological and clinical characteristics of 125 newborns born to women infected with COVID-19 by Turkish Neonatal Society. *Eur J Pediatr* 2020;180(3):733-742.
6. Salvatore CM, Han J-Y, Acker KP, et al. Neonatal management and outcomes during the COVID-19 pandemic: an observation cohort study. *Lancet Child Adolesc Health* 2020;4:721–727.
7. Gale C, Quigley MA, Placzek A, et al. Characteristics and outcomes of neonatal SARS-CoV-2 infection in the UK: a prospective national cohort study using active surveillance. *Lancet Child Adolesc Health* 2021;5:113–121.
8. Knight M, Bunch K, Vousden N, et al. Characteristics and outcomes of pregnant women admitted to hospital with confirmed SARS-CoV-2 infection in UK: national population based cohort study. *BMJ* 2020;369:m2107.

9. Olivini N, Calò Carducci FI, Santilli V, et al. A neonatal cluster of novel coronavirus disease 2019: clinical management and considerations. *Ital J Pediatr* 2020;46:180.
10. Ashary N, Bhide A, Chakraborty P, et al. Single-Cell RNA-seq Identifies Cell Subsets in Human Placenta That Highly Expresses Factors Driving Pathogenesis of SARS-CoV-2. *Front Cell Dev Biol* 2020;8:783.
11. Ovali F. SARS-CoV-2 Infection and the Newborn. *Front Pediatr* 2020;8:294–294.
12. Chamseddine RS, Wahbeh F, Chervenak F, et al. Pregnancy and Neonatal Outcomes in SARS-CoV-2 Infection: A Systematic Review. *J Pregnancy* 2020;2020:1-7.
13. Shende P, Gaikwad P, Gandhewar M, et al. Persistence of SARS-CoV-2 in the first trimester placenta leading to transplacental transmission and fetal demise from an asymptomatic mother. *Hum Reprod* 2021 Mar 18;36(4):899-906.
14. Fenizia C, Biasin M, Cetin I, et al. Analysis of SARS-CoV-2 vertical transmission during pregnancy. *Nat Commun* 2020;11:5128.
15. Facchetti F, Bugatti M, Drera E, et al. SARS-CoV2 vertical transmission with adverse effects on the newborn revealed through integrated immunohistochemical, electron microscopy and molecular analyses of Placenta. *EBioMedicine* 2020;59:102951.
16. Groß R, Conzelmann C, Müller JA, Stenger S, Steinhart K, Kirchhoff F, et al. Detection of SARS-CoV-2 in human breastmilk. *The Lancet* 2020;395:1757–1758.
17. Chambers C, Krogstad P, Bertrand K, et al. Evaluation for SARS-CoV-2 in Breast Milk From 18 Infected Women. *JAMA* 2020;324:1347–8.
18. Grimminck K, Santegoets LAM, Siemens FC, et al. No evidence of vertical transmission of SARS-CoV-2 after induction of labour in an immune-suppressed SARS-CoV-2-positive patient. *BMJ Case Rep* 2020;13:e235581.

19. Home - n.d. <https://pregcovid.com/> (27 May 2021, date last accessed).

20. Mahajan NN, Pednekar R, Patil SR, et al. Preparedness, administrative challenges for establishing obstetric services, and experience of delivering over 400 women at a tertiary care COVID-19 hospital in India. *Int J Gynecol & Obstet* 2020, 151: 188-196.

21. COVID-19 Neonatal guidelines by Kerala, NNF March 2020. <http://www.pedijatri.org/wp-content/uploads/2020/03/Neonatoski-vodic-iz-Kerale.pdf> (27 May 2021, date last accessed).

22. Chawla D, Chirla D, Dalwai S, et al. Perinatal-Neonatal Management of COVID-19 Infection - Guidelines of the Federation of Obstetric and Gynaecological Societies of India (FOGSI), National Neonatology Forum of India (NNF), and Indian Academy of Pediatrics (IAP). *Indian Pediatr* 2020;57:536–548.

23. World Health Organization. Frequently asked questions: breastfeeding and COVID-19: for health care workers, 12 May 2020. Geneva: World Health Organization; 2020.

24. Tiwari L, Taneja LN, Gupta S, et al. IAP ALS Update on Resuscitation Guidelines During COVID-19 Pandemic. *Indian J Pediatr* 2020:1–7.

25. Dong Y, Mo X, Hu Y, et al. Epidemiology of COVID-19 Among Children in China. *Pediatrics* 2020;145:e20200702.

26. Khalil A, Kalafat E, Benlioglu C, et al. SARS-CoV-2 infection in pregnancy: A systematic review and meta-analysis of clinical features and pregnancy outcomes. *EClinicalMedicine* 2020:100446.

27. Anand P, Yadav A, Debata P, et al. Clinical profile, viral load, management and outcome of neonates born to COVID 19 positive mothers: a tertiary care centre experience from India. *Eur J Pediatr* 2021;180:547–559.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
28. Chi H, Chiu NC, Tai YL, et al. Clinical features of neonates born to mothers with coronavirus disease-2019: A systematic review of 105 neonates. *J Microbiol Immunol Infect.* 2021;54(1):69-76.
 29. Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020;8:475–481.
 30. McLaren SH, Dayan PS, Fenster DB, et al. Novel Coronavirus Infection in Febrile Infants Aged 60 Days and Younger. *Pediatrics.* 2020;146(3):e20201550.
 31. Kotlyar AM, Grechukhina O, Chen A, et al. Vertical transmission of coronavirus disease 2019: a systematic review and meta-analysis. *Am J Obstet Gynecol* 2021;224:35-53.e3.
 32. Shah PS, Diambomba Y, Acharya G, Morris SK, Bitnun A. Classification system and case definition for SARS-CoV-2 infection in pregnant women, fetuses, and neonates. *Acta Obstet Gynecol Scand.* 2020;99(5):565-568. doi:10.1111/aogs.13870
 33. Zeng L, Xia S, Yuan W, et al. Neonatal Early-Onset Infection With SARS-CoV-2 in 33 Neonates Born to Mothers With COVID-19 in Wuhan, China. *JAMA Pediatr* 2020;174:722–5.
 34. Nanavati R, Mascarenhas D, Goyal M, et al. A single-center observational study on clinical features and outcomes of 21 SARS-CoV-2-infected neonates from India. *Eur J Pediatr* 2021;1–12.

Figure legends

Fig. 1. A course of neonates born to mothers with COVID-19.

*Roomed in with Mother: Exclusive breastfeeding (EBF) and skin-to-skin care contact was followed for all stable mother-infant dyads. EBF was practiced by counseling the mother to take utmost precautions by ensuring strict respiratory (wearing a medical mask) and hand hygiene before and after feeding or touching the baby. Unstable, preterm babies and babies born to sick mothers were given expressed breast milk (EBM) / pasteurized donor human milk (PDHM) from Human Milk Bank with feeding cup / spoon-feeding /or gastric tube feeding as per needs of the baby.

Fig. 2. Indications for NICU admissions in newborns of mothers with COVID-19.

Table 1: Details of the SARS-CoV-2 testing in neonates of mothers with COVID-19

Tests	Number	NICU Admission	Symptomatic	Asymptomatic	Discharged	Died
Only 1 st RT-PCR Positive	25	6	6*	19	23	2
Only 2 nd RT-PCR Positive	4	1	2	2	4	0
Both 1 st & 2 nd RT-PCR Positive	4	2	2	2	3	1
Total	33	9	10	23	30	3
<i>*One neonate in this category was managed in the postnatal wards for hyperbilirubinemia</i>						

Table 2: Demographic characteristics of neonates born to mothers with COVID-19

Characteristics	Total neonates n=524 (%)	SARS-CoV-2 infected n=33 (%)	SARS-CoV-2 non- infected* n=491 (%)	OR (95% CI)	<i>p</i> value
Male	281 (53.6)	13 (39.4)	268 (54.6)	0.54 (0.26-1.11)	0.090
Female	243 (46.4)	20 (60.6)	223 (45.4)		
Preterm delivery	25 (4.8)	3 (9.1)	22 (4.5)	2.13 (0.60-7.53)	0.203
Term delivery	499 (95.2)	30 (90.9)	469 (95.5)		
AGA	464 (88.5)	28 (84.8)	436 (88.8)	0.71 (0.26-1.91)	0.568
SGA	59 (11.3)	5 (15.2)	54 (11.0)		
Birthweight <1500 gm	6 (1.1)	1(3.0)	5 (1.0)	0.549	0.549
Birthweight 1500-2500 gm	118 (22.5)	8 (24.2)	110 (22.4)		
Birthweight >2500 gm	400 (76.3)	24 (72.7)	376 (76.6)		
AGA-Appropriate gestation age. SGA-Short for gestational age. OR-Odds ratio. CI-Confidence Interval.					
*Including a baby who died because of birth asphyxia and in whom nasopharyngeal swab collection was not possible.					

Table 3: Clinical characteristics and complications of neonates born to mothers with COVID-19

Characteristics	Total neonates n=523 (%)	SARS-CoV-2 infected n=33 (%)	SARS-CoV-2 non-infected* n=490 (%)	OR (95% CI)	<i>p</i> value
Complications in neonatal period	104 (19.9)	9 (27.3)	95 (19.4)	1.56 (0.70-3.46)	0.272
Congenital anomalies	5 (1.0)	0 (0)	5 (1.0)	1.32 (0.07-24.33)	
Birth asphyxia	7 (1.3)	1 (3.0)	6 (1.2)	2.52 (0.29-21.58)	0.368
Meconium aspiration syndrome	4 (0.8)	0 (0)	4 (0.8)	1.61(0.08-30.61)	
Hypocalcemia	3 (0.6)	1 (3.0)	2 (0.4)	7.62 (0.67-86.34)	0.178
Hypoglycemia	9 (1.7)	2 (6.1)	7 (1.4)	4.45 (0.89-22.33)	0.105
Sepsis	20 (3.8)	4 (12.1)	16 (3.3)	4.09 (1.28-13.00)	0.031
Hyperbilirubinemia	59 (11.3)	5 (15.2)	54 (11.0)	1.44 (0.53-3.89)	0.404

NICU-Neonatal Intensive Care Unit. OR-Odds ratio. CI-Confidence interval.

*Excluding a baby who died because of birth asphyxia and in whom nasopharyngeal swab collection was not possible.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Table 4: Clinical presentations in neonates born to mothers with COVID-19

Characteristics	Total neonates n=523 (%)	SARS-CoV-2 infected n=33 (%)	SARS-CoV-2 non-infected* n=490 (%)	OR (95% CI)	<i>p</i> value
Fever	4 (0.8)	1 (3.0)	3 (0.6)	5.07 (0.51-50.16)	0.230
Poor feeding	17 (3.3)	3 (9.1)	3 (0.6)	16.23 (3.14-83.87)	0.017
Respiratory distress	23 (4.4)	4 (12.1)	13 (2.7)	5.06 (1.55-16.49)	0.050
Oxygen saturation					
≤95% saturation	14 (2.7)	4 (12.1)	10 (2.0)	6.62 (1.95-22.39)	0.008
>95% saturation	509 (97.3)	29 (87.9)	480 (98.0)		

OR-Odds ratio. CI-Confidence interval.
*Excluding a baby who died because of birth asphyxia and in whom nasopharyngeal swab collection was not possible.

Table 5: Management of neonates born to mothers with COVID-19

Characteristics	Total neonates n=523 (%)	SARS-CoV-2 infected n=33 (%)	SARS-CoV-2 non-infected* n=490 (%)	OR (95% CI)	<i>p</i> value
Neonatal resuscitation	19 (3.6)	2 (6.1)	17 (3.5)	1.79 (0.39-8.12)	0.340
NICU admission	132 (25.2)	9 (27.3)	123 (25.1)	1.12 (0.51-2.47)	0.801
Bag and Mask	18 (3.4)	2 (6.1)	16 (3.3)	1.91 (0.42-8.68)	0.316
CPAP	15 (2.9)	4 (12.1)	11 (2.2)	6.00 (1.80-20.02)	0.011
Ventilator support	16 (3.1)	3 (9.1)	13 (2.7)	3.67 (0.99-13.58)	0.073
Treatments [†] (n=125)	125 (23.9)	9 (27.3)	116 (23.7)	1.21 (0.55-2.67)	0.639
Phototherapy	59 (11.3)	5 (15.1)	54 (11.0)		
Antibiotics	42 (8.0)	5 (15.1)	37 (7.6)		
Supportive therapy [‡]	66 (12.6)	3 (9.1)	63 (12.9)		

CPAP - Continuous positive airway pressure. OR-Odds ratio. CI-Confidence interval.

*Excluding a baby who died because of birth asphyxia and in whom nasopharyngeal swab collection was not possible.

[†]Neonates required either one or combinations of treatment.


[‡]Supportive therapy includes intravenous fluid, warmer care, measured feed, Haemo Gluco Test monitoring (HGT).

Table 6: Clinical and biochemical profile of nine neonates requiring NICU admission

Case	1	2	3	4	5	6	7	8	9
Hb gm/dl	15	15.8	16	7.6 (↓)	16.6	13.6	10.3(↓)	13	14.2
WBC (10 ³)	8.9	14.8	11.5	12.3	10.15	17.4	6.0	3.88 (↓)	13.4
Platelet Count (10 ⁵)	4.2	2.08	2.5	0.23 (↓)	2.92	4.1	0.22 (↓)	0.06 (↓)	2.65
LFT-Bilirubin/S GOT/SGPT	4.2/27/ 24	4/28/21	8.4/31/1 0	0.8/45/17	17/ 38/24	16.5 (Direct 6.8) 577/909	10.9 / 221 / 17	9.8/35/34	18/48/18
RFT-Na/ K/ BUN/Creatinine	146/5.0 21/0.8	141 /4.5 22 /0.9	144/5.6 15/0.5	135/ 4.8/27/- 0.7	138/4.3/52/ 0.7	129/5.5 44/0.5	134/4.6 16/1.1	143/4.86 8/0.8	136/4.7
Blood Culture	*	No Growth	*	No Growth	*	<i>Klebsiella Pneumonia</i>	<i>Klebsiella Pneumonia</i>	No Growth	33/0.7
X-ray Chest	Normal	Normal	Normal	Pneumonia	Normal	Normal	Normal	Normal	Normal
CRP	*	5	*	78(↑)	*	51(↑)	37(↑)	73(↑)	*
IL-6 / D-Dimer /LDH	*	*	*	**	*	IL-6, 140 (↑) LDH-371.5. D-dimer14.6	**	IL-6, 5000(↑)	*
CSF	*	*	*	**	*	Normal	**	**	*
Diagnosis	IDM	Birth Asphyxia	Hyperbi lirubine mia	PT-RDS- Sepsis-	Hyperbiliru binemia	Sepsis with Cholestatic Jaundice	PT-Sepsis – Succumbed	SGA- Hypoglycemia -Sepsis- Succumbed	Hyperbiliru binemia

* Not Indicated, **- Could not be done, ↑-increased, ↓-decreased,
In case numbers-1, 2, 3, 5, 9 these investigations were not indicated. In case numbers 4,7,8—CSF examination could not be done due to very poor general condition or low platelet counts, in case of number4,7,8, the tests could not be done because of logistic issues.
Hb- Hemoglobin, WBC-White blood count, LFT-Liver function test, SGOT- serum glutamic-oxaloacetic transaminase, SGPT- Serum glutamic-pyruvate transaminase, RFT-Renal function test, BUN-Blood Urea Nitrogen, CRP-C Reactive Protein, IL 6-Interleukin 6, LDH- Lactate dehydrogenase, CSF-Cerebrospinal Fluid. IDM-Infant of Diabetic Mother, PT-Preterm, RDS-Respiratory Distress Syndrome, SGA-Short for Gestational Age

Figure 1: Course of neonates born to mothers with COVID-19




Roomed in with Mother*: Exclusive breast feeding (EBF) and skin to skin care contact was followed for all stable mother-infant dyads. EBF was practiced by counselling the mother to take utmost precautions by ensuring strict respiratory (wearing a medical mask) and hand hygiene before and after feeding or touching the baby. Unstable, preterm babies and babies born to sick mothers were given expressed breast milk (EBM) / pasteurized donor human milk (PDHM) from Human Milk Bank with feeding cup / spoon feeding for gastric tube feeding as per needs of the baby.

Course of neonates born to mothers with COVID-19.

338x190mm (400 x 400 DPI)

Figure 2: Indications for NICU admissions in newborns of mothers with COVID-19



Indications for NICU admissions in newborns of mothers with COVID-19.

338x190mm (400 x 400 DPI)